Patient Alert Card

9064A1

Eliquis°

(apixaban)

Carry this card with you at all times

Show this card to your pharmacist, dentist and any other healthcare professionals that treat you.

Information for PATIENTS

- •Take Eliquis® regularly as instructed. If you miss a dose, take it as soon as you remember and continue to follow your dosing schedule.
- Do not stop taking Eliquis® without talking to your doctor, as you are at risk of suffering from a stroke or other complications.
- Eliquis® helps to thin your blood. However, this may increase your risk of bleeding.
- Signs and symptoms of bleeding include bruising or bleeding under the skin, tar-coloured stools, blood in urine, nose-bleed, dizziness, tiredness, paleness or weakness, sudden severe headache, coughing up blood or vomiting blood.
- If the bleeding does not stop on its own, immediately seek medical attention.
- If you need surgery, inform your doctor that you are taking Eliquis®.

I am under anticoagulation treatment with Eliquis® (apixaban) to prevent blood clots

This medicinal product is subject to additional monitoring. Patients and Health Care Professionals are asked to report any suspected adverse reactions via the national reporting system.

Information for HEALTHCARE PROFESSIONALS

- •Eliquis® (apixaban) is an oral anticoagulant acting by direct selective inhibition of factor Xa.
- •Eliquis® may increase the risk of bleeding. In case of major bleeding events, it should be stopped immediately.
- •Treatment with Eliquis® does not require routine monitoring of exposure. A calibrated quantitative anti-Factor Xa assay may be useful in exceptional situations, e.g., overdose and emergency surgery (prothrombin time (PT), international normalized ratio (INR) and activated partial thromboplastin time (aPTT) clotting tests are not recommended) see SmPC

Please complete this section or ask your doctor to do it

Name:	
Birth Date:	
Indication:	
Dose:	mg twice daily
Doctor's Name:	
Doctor's telephone:	
	_

