

Important Contact Details

My name:

My emergency contact (name and phone number):

Doctor's name – who prescribed Venclexta[®] (venetoclax):

Doctor's contact phone number:



Important Safety Information for **Healthcare Professionals** about Venclexta® (venetoclax) and TLS

This patient is taking Venclexta® (venetoclax) for CLL which may cause Tumour Lysis Syndrome (TLS) in the first 5 weeks of treatment.

- Evaluate blood chemistries: potassium, uric acid, phosphorus, calcium and creatinine.
- Urgently correct blood chemistry abnormalities. **This can be fatal, if left untreated.**
- If you suspect TLS, contact prescribing doctor listed on this card immediately.

Report side effects directly to AbbVie Biopharmaceuticals GmbH via Email: PV.MEA@abbvie.com, Hotline: 00966 55 828 2010 and National Pharmacovigilance Center Saudi Food and Drug Authority via SFDA Unified Call Center: 19999, E-mail: npc.drug@sfd.gov.sa, Website: <https://ade.sfda.gov.sa/>



This document is approved by The Executive Directorate of Pharmacovigilance, at SFDA

Patient Card

Safety Information about Venclexta[®] (venetoclax) for Patients with Chronic Lymphocytic Leukaemia (CLL)

Venclexta[®] (venetoclax) is used to treat patients with CLL. This card contains information you need to know about Tumour Lysis Syndrome and Venclexta[®] (venetoclax).



- Keep this patient card with you all the time.
- Show it to any doctor involved in your medical care or if you go to the emergency department.
- Always take Venclexta[®] (venetoclax) exactly as your doctor has told you to.

What should I know about TLS?

During your treatment, you may develop high levels of certain body salts (such as potassium and uric acid) in the blood. This can lead to a condition called Tumour Lysis Syndrome (TLS) within the first 5 weeks of treatment.

TLS symptoms:

- fever or chills
- nausea or vomiting
- feeling confused
- feeling short of breath
- irregular heart beat
- dark or cloudy urine
- feeling unusually tired
- muscle pain or joint pain
- fits or seizures
- abdominal pain and bloating

What should I do if I get TLS symptoms?

If you experience any TLS symptoms during your treatment, you should:



Call your doctor or seek medical attention immediately, and



Stop taking Venclexta[®] (venetoclax)

Read the Patient Information Leaflet (PIL) for more information.

How can I reduce my risk of getting TLS?



Drink at least **1.5 to 2 liters of water daily**:

- starting from 2 days before your first dose, and
- on the day of your first dose, and
- every day during first 5 weeks of treatment.

Drinking plenty of water will help to remove high levels of blood salts through your urine.



Attend your scheduled **blood test appointments**, as they will help your doctor identify TLS.



Before your treatment with Venclexta[®]

(venetoclax), start another medicine that helps prevent the build-up of the blood salt (uric acid) in your body, as instructed by your doctor.