



Apixaban 2.5mg & 5mg

Prescriber Guide

This Prescriber Guide is not a substitute for the **Pixar®** Summary of Product Characteristics (SmPC). Please consult the SmPC for full prescribing information.

This document has been reviewed and approved by Saudi Food and Drug Authority (SFDA)

This educational material is provided to further minimize the risk of bleeding that is associated with the use of apixaban and to guide healthcare professionals in managing that risk.

Table of Contents

Patient Alert Card	3
Therapeutic indication: Prevention of stroke and systemic embolism in adult patients with non-valvular atrial fibrillation (NVAf) with one or more risk factors	
Dosing recommendations	4
Dose reduction	4
Missed dose	5
Patients with renal impairment	5
Patients with hepatic impairment	6
Patients undergoing catheter ablation	6
Patients undergoing cardioversion	6
Therapeutic indication: Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrent DVT and PE in adults	
Dosing recommendations	7
Missed dose	8
Patients with renal impairment	8
Patients with hepatic impairment	8
Hemodynamically unstable PE patients or patients who require thrombolysis or pulmonary embolectomy	9
Patients with active cancer	9
Therapeutic indication: Prevention of venous thromboembolic events (VTE) in adult patients who have undergone elective hip or knee replacement surgery	
Dosing recommendations	10
Missed dose	10
Patients with renal impairment	10
Patients with hepatic impairment	11
Switching to and from apixaban	12
Populations potentially at higher risk of bleeding	13
Surgery and invasive procedures	15
Temporary discontinuation	15
Spinal/epidural anaesthesia or puncture	15
Management of overdose and haemorrhage	17
Use of coagulation testsReferences	17

Patient Alert Card

A Patient Alert Card must be provided to each patient who is prescribed apixaban 2.5 mg or 5 mg, and the importance and consequences of anticoagulant therapy should be explained.

Specifically, the prescriber should talk to patients about the importance of treatment compliance, the signs or symptoms of bleeding, and when to seek attention from a healthcare professional.

This Patient Alert Card provides information to healthcare professionals on the anticoagulant therapy and contains important contact information in the event of emergencies.

Patients should always be advised to carry the Patient Alert Card with them and to show it to every healthcare professional. They should also be reminded about the need to inform healthcare professionals that they are taking apixaban if they require surgery or invasive procedures.

Additional copies of the Patient Alert Card can be obtained by contacting Sudair Pharma Company via pharmacovigilance@sudairpharma.com,

phone: +966-920001432 ext.107

Therapeutic indication: Prevention of stroke and systemic embolism in adult patients with non-valvular atrial fibrillation (NVAF) with one or more risk factors

Risk factors for stroke in NVAF include prior stroke or transient ischemic attack, age ≥ 75 years, hypertension, diabetes mellitus, and symptomatic heart failure (NYHA Class $\geq II$).

Dosing recommendations

The recommended dose of apixaban is 5 mg taken orally twice daily with water, with or without food. Therapy should be continued long-term (Figure 1).

Figure 1




For patients who are unable to swallow whole tablets, apixaban tablets may be crushed and suspended in water, or 5% dextrose in water (D5W), or apple juice or mixed with apple puree and immediately administered orally.

Alternatively, apixaban tablets may be crushed and suspended in 60 mL of water or D5W and immediately delivered through a nasogastric tube. Crushed apixaban tablets are stable in water, D5W, apple juice, and apple puree for up to 4 hours.

Dose reduction

In patients with at least two of the following characteristics: age ≥ 80 years, body weight ≤ 60 kg, or serum creatinine ≥ 1.5 mg/dL (133 μ mol/L), the recommended dose of apixaban is 2.5 mg taken orally twice daily (Figure 2).

Patients with exclusive criteria of severe renal impairment (creatinine clearance [CrCl] 15–29 mL/min) should also receive apixaban 2.5 mg twice daily (Figure 2).

Figure 2


Missed dose

If a dose is missed, the patient should take apixaban immediately and then continue with twice daily intake as before.

Patients with renal impairment

Renal impairment	
Dialysis	Not recommended
Renal failure (CrCl <15 ml/min)	Not recommended
Severe renal impairment (CrCl 15–29 ml/min)	Dose reduction to 2.5 mg twice daily
Mild (CrCl 51–80 ml/min) or moderate (CrCl 30–50 ml/min) renal impairment	5 mg twice daily. No dose adjustment required unless the patient fulfils criteria for dose reduction to 2.5 mg twice daily based on age, body weight and/or serum creatinine (refer to dosing section)

Patients with hepatic impairment

Epatic impairment

Hepatic disease associatedwith coagulopa- thy and clinically relevant bleeding risk	Contraindicated
Severe hepatic impairment	Not recommended
Mild or moderate hepatic impairment (Child Pugh A or B)	Use with caution No dose adjustment required

Prior to initiating apixaban, liver function testing shouldbe performed. Patients with elevated liver enzymes alanine aminotransferase (ALT)/aspartate aminotransferase (AST) >2 x ULNorto-
t bilirubin ≥1.5 x ULNwere excluded in clinical trials. Therefore, apixaban should be used cau-
tiously in this population.

Patients undergoing catheter ablation

Apixaban can be continued in patients undergoing catheter ablation for atrial fibrillation.

Patients undergoing cardioversion

Apixaban can be initiated or continued in NVAF patients who may require cardioversion.

For patients not previouslytreatedwith anticoagulants, exclusion of left atrial thrombususing
an image-guided approach (e.g. transesophageal echocardiography [TEE] or computed tomo-
graphic scan [CT]) prior to cardio- version should be considered, in accordance with established
medical guidelines. For patients in whom a prior thrombus has been detected, established med-
ical guidelines should be followed prior to cardioversion.

Patient status	Patient qualifies for dose reduction?	Dosing regimen
Initiating treatment with apixaban	No	5 mg twice daily for at least 2.5 days (5 single doses) before cardioversion
	Yes	2.5 mg twice daily for at least 2.5 days (5 single doses) before cardioversion
Insufficient time prior to cardioversion to adminis- ter 5 doses of apixaban	No	10 mg loading dose at least 2 hours before cardioversion, followed by 5 mg twice daily
	Yes	5 mg loading dose at least 2 hours before cardioversion, followed by 2.5 mg twice daily

For all patients undergoing cardioversion, confirmation should be sought prior to cardioversion
that the patient has taken apixaban as prescribed. Decisions on initiation and duration of treat-
ment should take established guideline recommendations for anticoagulant treatment in pa-
tientsundergoing cardioversion into account.

Therapeutic indication: Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrent DVT and PE in adults

Dosing recommendations









The recommended dose of apixaban for the treatment of acute DVT and treatment of PE is 10 mg taken orally twice daily for the first 7 days followed by 5 mg taken orally twice daily with water, with or without food.

As per available medical guidelines, short duration of treatment (at least 3 months) should be based on major transient/reversible risk factors (e.g. recent surgery, trauma, immobilization).

The recommended dose of apixaban for the prevention of recurrent DVT and PE is 2.5 mg taken orally twice daily with water, with or without food.

When prevention of recurrent DVT and PE is indicated, the 2.5 mg twice daily dose should be initiated following completion of 6 months of treatment with apixaban 5 mg twice daily or with another anticoagulant, as indicated in Figure 3.

Figure 3

Dosing Schedule	 Morning	 Night	Daily Dose
Day 1 – 7: 10 mg twice daily →	 5 mg 5 mg	 5 mg 5 mg	20 mg
Day 8 onwards: 5 mg twice daily →	 5 mg	 5 mg	10 mg
Prevention of recurrent DVT and/or PE following completion of 6 months anticoagulation treatment			
2.5 mg twice daily →	 2.5 mg	 2.5 mg	5 mg

The duration of overall therapy should be individualized after careful assessment of the treatment benefit against the risk for bleeding.

For patients who are unable to swallow whole tablets, apixaban tablets may be crushed and suspended in water, or 5% dextrose in water (D5W), or apple juice or mixed with apple puree and immediately administered

orally. Alternatively, apixaban tablets may be crushed and suspended in 60 mL of water or D5W and immediately delivered through a nasogastric tube. Crushed apixaban tablets are stable in water, D5W, apple juice, and apple puree for up to 4 hours.

Missed dose

If a dose is missed, the patient should take apixaban immediately and then continue with twice daily intake as before.

Patients with renal impairment

Renal impairment	
Dialysis	Not recommended
Renal failure (CrCl <15 ml/min)	Not recommended
Severe renal impairment (CrCl 15–29 ml/min)	Use with caution
Mild (CrCl 80–51 ml/min) or moderate (CrCl 50–30 ml/min) renal impairment	No dose adjustment

Patients with hepatic impairment

Renal impairment	
Hepatic disease associated with coagulopathy and clinically relevant bleeding risk	Contraindicated
Severe hepatic impairment	Not recommended
Mild or moderate hepatic impairment (Child Pugh A or B)	Use with caution No dose adjustment required

Prior to initiating apixaban, liver function testing should be performed. Patients with elevated liver enzymes ALT/AST >2 x ULN or total bilirubin ≥1.5 x ULN were excluded in clinical trials. Therefore, apixaban should be used cautiously in this population.

Hemodynamically unstable PE patients or patients who require thrombolysis or pulmonary embolectomy

Apixaban is not recommended as an alternative to unfractionated heparin in patients with PE who are hemodynamically unstable or may receive thrombolysis or pulmonary embolectomy.

Patients with active cancer

Efficacy and safety of apixaban in the treatment of DVT, treatment of PE, and prevention of recurrent DVT and PE in patients with active cancer have not been established.

Therapeutic indication: Prevention of venous thromboembolic events (VTE) in adult patients who have undergone elective hip or knee replacement surgery

Dosing recommendations

The recommended dose of apixaban is 2.5 mg taken orally twice daily with water, with or without food. The initial dose should be taken 12 to 24 hours after surgery.

Physicians may consider the potential benefits of earlier anticoagulation for VTE prophylaxis as well as the risks of post-surgical bleeding in deciding on the time of administration within this time window.

In patients undergoing **hip replacement surgery** the recommended duration of treatment is **32 to 38 days**.

In patients undergoing **knee replacement surgery** the recommended duration of treatment is **10 to 14 days**.

For patients who are unable to swallow whole tablets, apixaban tablets may be crushed and suspended in water, or 5% dextrose in water (D5W), or apple juice or mixed with apple puree and immediately administered

orally. Alternatively, apixaban tablets may be crushed and suspended in 60 mL of water or D5W and immediately delivered through a nasogastric tube. Crushed apixaban tablets are stable in water, D5W, apple juice, and apple puree for up to 4 hours.

Missed dose

If a dose is missed, the patient should take apixaban immediately and then continue with twice daily intake as before.

Patients with renal impairment

Renal impairment	
Dialysis	Not recommended
Renal failure (CrCl <15 ml/min)	Not recommended
Severe renal impairment (CrCl 15–29 ml/min)	Use with caution
Mild (CrCl 30–50 ml/min) or moderate (CrCl 15–29 ml/min) renal impairment	No dose adjustment required

Patients with hepatic impairment
Hepatic impairment

Hepatic disease associated with coagulopathy and clinically relevant bleeding risk	Contraindicated
Severe hepatic impairment	Not recommended
Mild or moderate hepatic impairment (Child Pugh A or B)	Use with caution No dose adjustment required

Prior to initiating apixaban, liver function testing should be performed. Patients with elevated liver enzymes ALT/AST >2 x ULN or total bilirubin ≥1.5 x ULN were excluded in clinical trials. Therefore, apixaban should be used cautiously in this population.

Switching to and from Apixaban


Switching treatment from parenteral anticoagulantst to apixaban and vice versa) can be done at the next scheduled dose.

These agents should not be administered simultaneously.

Switching from vitamin K antagonist (VKA) therapy to apixaban

When converting patients from VKA therapy to apixaban, discontinue warfarin or other VKA therapy and start apixaban when the international normalized ratio (INR) is <2.0 (Figure 4).

Figure 4



Switching from apixaban to VKA therapy

When converting patients from apixaban to VKA therapy, continue administration of apixaban for at least 2 days after beginning VKA therapy. After 2 days of co-administration of apixaban with VKA therapy, obtain an INR prior to the next scheduled dose of apixaban. Continue co-administration of apixaban and VKA therapy until the INR is ≥ 2.0 .

Populations potentially at higher risk of bleeding

Several sub-groups of patients are at increased risk of bleeding and should be carefully monitored for signs and symptoms of bleeding complications. Apixaban should be used with caution in conditions with an increased hemorrhagic risk. Apixaban administration should be discontinued if severe hemorrhage occurs.

Lesion or condition if considered a significant risk factor for major bleeding and where use is contraindicated

This includes:

- Active clinically significant bleeding
- Hepatic disease associated with coagulopathy and clinically relevant bleeding risk
- Current or recent gastrointestinal ulceration
- Presence of malignant neoplasms at high risk of bleeding
- Recent brain or spinal injury
- Recent brain, spinal or ophthalmic surgery
- Recent intracranial hemorrhage
- Known or suspected esophageal varices, arteriovenous malformations, vascular aneurysms or major intraspinal or intracerebral vascular abnormalities

Interactions with other medicinal products affecting haemostasis

Anticoagulants

- Unfractionated heparin (UFH), low molecular weight heparins (e.g. enoxaparin, dalteparin), heparin derivatives (e.g. fondaparinux)
- Oral anticoagulants (e.g. warfarin, rivaroxaban, dabigatran)

Due to an increased bleeding risk, concomitant treatment with apixaban and any other anticoagulant agent is **contraindicated**, except under specific circumstances of switching anticoagulant therapy, when UFH is given at doses necessary to maintain an open central venous or arterial catheter, or when UFH is given during catheter ablation for atrial fibrillation

Platelet aggregation inhibitors, SSRIs/ SNRIs and NSAIDs

- Unfractionated heparin (UFH), low molecular weight heparins (e.g. enoxaparin, dalteparin), heparin derivatives (e.g. fondaparinux)
- Selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs)
- Acetylsalicylic acid (ASA)
- Non-steroidal anti-inflammatory drugs (NSAIDs)

The concomitant use of apixaban with antiplatelet agents increases the risk of bleeding. Care is to be taken if patients are treated concomitantly with SSRIs/SNRIs or NSAIDs, including ASA

- Medicinal products associated with serious bleeding are not recommended concomitantly with apixaban, such as: thrombolytic agents, GPIIb/IIIa receptor antagonists, thienopyridines (e.g. clopidogrel), dipyridamole, dextran and sulfinpyrazone.

Interactions with other medicinal products affecting haemostasis

Renal impairment	<p>See sections on patients with renal impairment under dosing recommendations for each separate indication</p> <ul style="list-style-type: none"> Use is not recommended in patients with CrCl <15 ml/min or patients undergoing dialysis No dose adjustment is required in patients with mild or moderate renal impairment <p>Patients with NVAf</p> <ul style="list-style-type: none"> Patients with severe renal impairment (CrCl 15–29 ml/min) should receive the lower dose of apixaban 2.5 mg twice daily Patients with serum creatinine ≥1.5 mg/dL (133 μmol/L) associated with age ≥80 years or body weight ≤60 kg should receive the lower dose of apixaban 2.5 mg twice daily
Elderly	<ul style="list-style-type: none"> No dose adjustment required <p>Patients with NVAf</p> <ul style="list-style-type: none"> No dose adjustment required except in combination with other factors
Low body weight ≤60 kg	<ul style="list-style-type: none"> No dose adjustment required <p>Patients with NVAf</p> <ul style="list-style-type: none"> No dose adjustment required except in combination with other factors
Concomitant use with strong inhibitors of both CYP3A4 and P-gp	<ul style="list-style-type: none"> Apixaban is not recommended in patients receiving concomitant systemic treatment with strong inhibitors of both CYP3A4 and P-gp, such as azole-antimycotics (e.g. ketoconazole, itraconazole, voriconazole and posaconazole) and HIV protease inhibitors (e.g. ritonavir)
Concomitant use with agents not considered strong inhibitors of both CYP3A4 and P-gp	<ul style="list-style-type: none"> No dose adjustment for apixaban is required when co-administered with, for example, diltiazem, naproxen, clarithromycin, amiodarone, verapamil and quinidine
Concomitant use with strong inducers of both CYP3A4 and P-gp	<ul style="list-style-type: none"> The concomitant use of apixaban with strong inducers of both CYP3A4 and P-gp (e.g. rifampicin, phenytoin, carbamazepine, phenobarbital or St. John's Wort) may lead to a ~50% reduction in apixaban exposure and should be used with caution <p>Treatment of DVT or PE</p> <ul style="list-style-type: none"> Apixaban is not recommended

Surgery and invasive procedures

Apixaban should be discontinued prior to elective surgery or invasive procedures (excluding cardioversion or catheter ablation) with a risk of bleeding (see *table below*).

If surgery or invasive procedures cannot be delayed, exercise appropriate caution, taking into consideration an increased risk of bleeding. This risk of bleeding should be weighed against the urgency of intervention.

In the event a patient treated with apixaban requires an elective procedure, such as surgery or an invasive procedure associated with an increased risk of bleeding, apixaban should be discontinued for a sufficient period of time prior to the procedure to reduce the risk of anticoagulant-related bleeding. The half-life of apixaban is approximately 12 hours. Given that apixaban is a reversible factor Xa inhibitor, its anticoagulant activity should abate within 24 to 48 hours from the last administered dose.

Discontinuation of apixaban prior to elective surgery / invasive procedure

Low risk of bleeding (includes interventions for which bleeding, if it occurs, will be minimal, non-critical in its location and/or easily controlled by simple mechanical hemostasis)	At least 24 hours prior to elective surgery or invasive procedure
Moderate or high risk of bleeding (includes interventions for which the probability of clinically significant bleeding cannot be excluded, or for which the risk of bleeding would be unacceptable)	At least 48 hours prior to elective surgery or invasive procedure

Temporary discontinuation

Discontinuing anticoagulants, including apixaban, for active bleeding, elective surgery, or invasive procedures places patients at an increased risk of thrombosis. Lapses in therapy should be avoided and if anticoagulation with apixaban must be temporarily discontinued for any reason, therapy should be restarted as soon as possible provided the clinical situation allows and adequate hemostasis has been established.

Spinal / epidural anesthesia or puncture

When neuraxial anesthesia (spinal/epidural anesthesia) or spinal/epidural puncture is employed, patients treated with antithrombotic agents for prevention of thromboembolic complications are at risk of developing an epidural or spinal hematoma, which can result in long-term or permanent paralysis. Post-operative indwelling epidural or intrathecal catheters must be removed **at least 5 hours** prior to the first dose of apixaban.


Guidance on the use of apixaban in patients with indwelling intrathecal or epidural catheters

There is no clinical experience with the use of apixaban with indwelling intrathecal or epidural catheters. In case there is such need and based on the general pharmacokinetic characteristics of apixaban, a time interval of **20 to 30 hours** (i.e. 2 x half-life) between the last dose of apixaban and catheter withdrawal should elapse, and at least one dose should be omitted before catheter withdrawal. The next dose of apixaban may be given **at least 5 hours** after catheter removal. As with all anticoagulant drugs, experience with

neuraxial blockade is limited and extreme caution is, therefore, recommended when using apixaban in the presence of neuraxial blockade (Figure 5).

Patients are to be frequently monitored for signs and symptoms of neurological impairment (e.g. numbness or weakness of the legs, bowel or bladder dysfunction). If neurological compromise is noted, urgent diagnosis and treatment are necessary.

Figure 5



Management of overdose and hemorrhage

Overdose of apixaban may result in a higher risk of bleeding. In the event of hemorrhagic complications, treatment must be discontinued, and the source of bleeding investigated. The initiation of appropriate treatment, e.g. surgical hemostasis, the transfusion of fresh frozen plasma, or the administration of a reversal agent for factor Xa inhibitors should be considered.

In controlled clinical trials, orally administered apixaban in healthy subjects at doses up to 50 mg daily for 3 to 7 days (25 mg twice daily for 7 days or 50 mg once daily for 3 days) had no clinically relevant adverse effects.

In healthy subjects, administration of activated charcoal 2 and 6 hours after ingestion of a 20 mg dose of apixaban reduced mean AUC by 50% and 27%, respectively, and had no impact on C_{max}. Mean half-life decreased from 13.4 hours when apixaban was administered alone to 5.3 hours and 4.9 hours, respectively, when activated charcoal was administered 2 and 6 hours after apixaban. Thus, administration of activated charcoal may be useful in the management of overdose or accidental ingestion.

For situations when reversal of anticoagulation is needed due to life-threatening or uncontrolled bleeding, a reversal agent for factor Xa inhibitors is available. Administration of prothrombin complex concentrates (PCCs) or recombinant factor VIIa may also be considered. Reversal of apixaban pharmacodynamic effects, as demonstrated by changes in the thrombin generation assay, was evident at the end of infusion and reached baseline values within 4 hours after the start of a 4-factor PCC 30-minute infusion in healthy subjects. However, there is no clinical experience with the use of 4-factor PCC products to reverse bleeding in individuals who have received apixaban. Currently there is no experience with the use of recombinant factor VIIa in individuals receiving apixaban. Re-dosing of recombinant factor VIIa could be considered and titrated depending on improvement of bleeding.

Depending on local availability, consultation of a coagulation expert should be considered in case of major bleeding.

Hemodialysis decreased AUC by 14% in subjects with end-stage renal disease, when a single dose of apixaban 5 mg was administered orally. Therefore, hemodialysis is unlikely to be an effective means of managing apixaban overdose.

Use of coagulation tests

Routine clinical monitoring is not required with apixaban treatment. However, a calibrated quantitative anti-Factor Xa assay may be useful in exceptional situations where knowledge of apixaban exposure may help to inform clinical decisions, e.g. overdose and emergency surgery.

Prothrombin time (PT), INR and activated partial thromboplastin time (aPTT)

Changes observed in these clotting tests at the expected therapeutic dose are small and subject to a high degree of variability. They are not recommended to assess the pharmacodynamic effects of apixaban.

In the thrombin generation assay, apixaban reduced endogenous thrombin potential, a measure of thrombin generation in human plasma.

Anti-FXa assays

Apixaban also demonstrates anti-FXa activity as evident by reduction in FXa enzyme activity in multiple commercial anti-FXa kits; however, results differ across kits. Data from clinical trials are only available for the **Pixar®** Heparin chromogenic assay. Anti-FXa activity exhibits a close direct linear relationship with apixaban plasma concentration, reaching maximum values at the time of apixaban peak plasma concentrations.

The relationship between apixaban plasma concentration and anti-FXa activity is approximately linear over a wide dose range of apixaban.

Table 1 shows the predicted steady-state exposure and anti-FXa activity for each indication. In patients taking apixaban for the prevention of VTE following hip or knee replacement surgery, the results demonstrate a less than 1.6-fold fluctuation in peak-to-trough levels. In NVAf patients taking apixaban for the prevention of stroke and systemic embolism, the results demonstrate a less than 1.7-fold fluctuation in peak-to-trough levels. In patients taking apixaban for the treatment of DVT and PE or prevention of recurrent DVT and PE, the results demonstrate a less than 2.2-fold fluctuation in peak-to-trough levels.

Predicted apixaban steady-state exposure and anti-FXa activity				
	Apixaban C _{max} (ng/mL)	Apixaban C _{min} (ng/mL)	anti-FXa activity max (IU/mL)	anti-FXa activity min (IU/mL)
	Median [5th, 95th percentile]			
Prevention of VTE: elective hip or knee replacement surgery				
2.5 mg twice daily	146 ,41] 77]	109 ,23] 51]	2.4 ,0.67] 1.3]	1.8 ,0.37] 0.84]
Prevention of stroke and systemic embolism: NVAF				
2.5 mg twice daily*	221 ,69] 123]	162 ,34] 79]	3.3 ,1.0] 1.8]	2.4 ,0.51] 1.2]
5 mg twice daily	321 ,91] 171]	230 ,41] 103]	4.8 ,1.4] 2.6]	3.4 ,0.61] 1.5]
Treatment of DVT, treatment of PE and prevention of recurrent DVT and PE				
2.5 mg twice daily	153 ,30] 67]	90 ,11] 32]	2.5 ,0.46] 1.0]	1.4 ,0.17] 0.49]
5 mg twice daily	302 ,59] 132]	177 ,22] 63]	5.2 ,0.91] 2.1]	2.9 ,0.33] 1.0]
10 mg twice daily	572 ,111] 251]	335 ,41] 120]	10.8 ,1.8] 4.2]	5.8 ,0.64] 1.9]

* Dose adjusted population based on at least 2 of 3 dose reduction criteria as shown in Figure 2

Reporting adverse reactions:

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Reporting forms and information can be found at SFDA as following:

Website: <https://ade.sfda.gov.sa>

Email: npc.drug@sfda.gov.sa

Call center: 19999



Also, you can report to Sudair Pharma Company via:

Phone: +966-920001432 Ext. 107

Email: Pharmacovigilance@SudairPharma.com

Mobile: +966 546030507



References

1. Reference product in EMC website: <https://www.medicines.org.uk/emc/>
2. Sudair Pharma Company. **Pixar®** 5mg film coated tablets Summary of Product Characteristics.
3. Sudair Pharma Company. **Pixar®** 2.5mg film coated tablets Summary of Product Characteristics.