


# Borvix® (Bortezomib) 3.5mg powder for solution for injection

## INDUCTION REGIMES PRIOR TO TRANSPLANT:

Dosing and duration of treatment



\*Dexamethasone is administered orally at 40 mg on days 1, 2, 3, 4 and 8, 9, 10, 11 of the Bortezomib treatment cycles.

\*\*Thalidomide dose is increased to 100 mg from week 3 of cycle 1 only if 50 mg is tolerated and to 200 mg from cycle 2 onwards if 100 mg is tolerated.

**Patients receiving Bortezomib in combination with Thalidomide should adhere to the pregnancy prevention programme of Thalidomide. Refer to the SmPC for additional information.**

\*\*\*Up to 6 cycles may be given to patients who achieve at least a partial response after 4 cycles.

### References

1. Borvix® SmPC and leaflet
2. Transplant Induction Regimens Graph. Retrieved 9 July 2020, from <https://www.medicines.org.uk/emc/rmm/1429/Document>

**To Report any Adverse Drug Reaction or Concerns Please call:  
Pharmacovigilance Department at MS Pharma:**

- Email: [pharmacovigilance@mspharma.com](mailto:pharmacovigilance@mspharma.com)
- Website: [www.mspharma.com](http://www.mspharma.com)
- Phone No: + 966112790122 Ext. 6013

**The National Pharmacovigilance Center (NPC): (Saudi food and drug authority)**

- Email: [npc.drug@sfd.gov.sa](mailto:npc.drug@sfd.gov.sa)
- Call Center: 19999
- Website: <https://ade.sfd.gov.sa/>
- QR Code:

