


i-SECURE

Imnovid® (Pomalidomide)

PRODUCT HANDLING INSTRUCTIONS FOR IMMUNOMODULATORY AGENTS

Points to consider for handling the medicinal product: for healthcare professionals, patients, family members, and caregivers:

- If you are a woman who is pregnant or suspect that you may be pregnant, you should not handle the blister or capsule.
- Wear disposable gloves when handling product and/or packaging (ie, blister or capsule).
- Capsules can occasionally become damaged when pressing them out of the blister, especially when the pressure is put onto the middle of the capsule. The pressure should be located at one site only, which reduces the risk of the capsule deforming or breaking (see figure below).
- Healthcare professionals, family members, and caregivers should wear disposable gloves when handling the blister or capsule.
- Use proper technique when removing gloves to prevent potential skin exposure (see picture below).
- Place gloves in a sealable plastic polyethylene bag, wash hands and dispose them according to local requirements.
- Wash hands thoroughly with soap and water after removing gloves.
- Patients should be advised never to give medication to another person.



If a drug product package looks visibly damaged, use the following extra precautions to prevent exposure:

- If the outer carton is visibly damaged – **do not open**.
- If blister strips are damaged or leaking or capsules look damaged or leaking – **close the outer carton immediately**.
- Place the product inside a sealable plastic polyethylene bag.
- Dispose of any unused medication according to local regulations.


If product is released or spilled, take proper precautions to minimize exposure by using appropriate personal protection:

- If capsules are crushed or broken, dust containing drug substance may be released. Avoid dispersing the powder and avoid breathing on or inhaling the powder.
- Wear disposable gloves to clean up the powder.
- Place a damp cloth or towel over the powder area to minimize entry of powder into the air. Add excess liquid to allow the material to enter solution. After handling, clean the area thoroughly with soap and water, then dry it.
- Place all contaminated materials, including damp cloth or towel, and the gloves, into a sealable polyethylene plastic bag. Dispose of it according to local requirements for medicinal products.
- Wash your hands thoroughly with soap and water after removing the gloves.
- Please report the release/spillage in accordance with local regulations or to the healthcare professional immediately.

If the contents of the capsule are attached to the skin or mucous membranes

- If you touch the drug powder, please wash exposed area thoroughly with running water and soap.
- If drug powder makes contact with one or both of your eyes, remove and discard any contact lenses in use. Then, thoroughly flush eyes with water for at least 15 minutes. If irritation occurs, please contact an ophthalmologist.

Proper Technique for Removing Gloves



- Grasp the outside edge near wrist (1).
- Peel away from hand, turning glove inside-out (2).
- Hold in opposite gloved hand (3).
- Slide ungloved finger under the wrist of the remaining glove, being careful not to touch the outside of the glove (4).
- Peel off from inside, creating a bag for both gloves.
- Discard in appropriate container.
- Wash your hands with soap and water thoroughly.