

Methotrexate BOS

Methotrexate 2.5 mg Tablet

Patient Card

Medication Errors

Oral methotrexate medicines used for conditions requiring once weekly treatment

This patient card is only intended for patients who use a methotrexate-containing medicine for indications: rheumatoid arthritis, juvenile idiopathic arthritis (JIA), psoriasis and lymphatic leukaemia as per indications listed in the package leaflet.

If you use methotrexate for one of the above-mentioned indications, you should only take methotrexate once a week.

Day of the Week for Intake

Patient Name

Patient Dose

Physician Name

Physician Phone Number

Do not take more than the prescribed dose.

Overdose could lead to serious adverse effects and may be fatal.
Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhea, vomiting, skin rashes, bleeding or unusual weakness.

If you think you have taken more than the prescribed dose, consult a physician immediately.

Always show this card to health care professionals not familiar with your methotrexate treatment to alert them about your once weekly use (e.g. on hospital admission, change of care).

For more information, please read the patient leaflet inserted in the package.

You can report any side effects to SFDA via:

The National Pharmacovigilance Centre (NPC)

SFDA Call Centre: 19999 | E-mail: npc.drug@sfd.gov.sa
Website: <https://ade.sfd.gov.sa/>

**Or to the pharmacovigilance department
of Boston Oncology Arabia Limited via:**

Boston Oncology Arabia Limited

Airport Road, 13413, Riyadh, Saudi Arabia
Phone: +966 54 764 3672 | E-mail: pv@bostononcology.com

**By reporting side effects, you can help provide more information on
the safety of this medicine.**

