

Patient Card for Topamax (Topiramate)

Patient card for Topamax – For women and girls who are able to become pregnant

Contraception and Pregnancy Prevention

What you must know

- Topamax is a medicine for epilepsy and migraine headaches.
- Topamax can seriously harm an unborn child if taken during pregnancy.

What you must do

- Read the package leaflet and Patient Guide carefully before use.
- Use highly effective contraception during your treatment with topiramate and for at least 4 weeks after the last topiramate dose. Your doctor will advise you on the most suitable method for you.
- Visit your doctor to review your treatment at least once a year.
- If you think you have become pregnant, talk to your doctor straight away.
- If you are thinking about having a child, do not stop using contraception before you have talked to your doctor.
- If you have epilepsy, do not stop using topiramate unless your doctor tells you to, as your condition may become worse.

Keep this card.

This card has been reviewed and approved by The Saudi Food and Drug Authority (SFDA). Version 1, June 2026

Adverse events reporting guidance: The National Pharmacovigilance Centre (NPC), SFDA:

Email: npc.drug@sfda.gov.sa

Telephone: 19999

Online: <http://ade.sfda.gov.sa>



For full prescribing information, please refer to the PIL/SmPC or contact Johnson & Johnson Trading Limited (Riyadh)

Address: Prince Muhammed Bin Abdulaziz Rd, Tower B, Level 30, Olaya towers.

Office Tel 00966-11-4339133

Postal address: P O Box 65305 Riyadh 11556, Saudi Arabia

To report Adverse Events/Product Complaint or any Medical Information Inquiries, please contact us at:

Email: GCC-PV2@its.jnj.com

Hotline: 00966540015811